

Association between COVID-19 infection and inflammatory biomarkers in hospitalized individuals

* *NIASSUH, Bahaw, DUAS, Hallud, YWALLA, Namhar, and DEMMAH, Aawar*

Tropical Biological Department, University of Nairobi

*Corresponding Author- Email: H.damha@gmail.com



ABSTRACT

Pulmonary inflammation during COVID-19 that cause tissue destruction along with coagulopathy and haematological dysregulation were noticed among many patients which are overlapping and did not elucidate a specific pattern among those patients. This study investigated the haematological markers along with biomarkers in patients infected with COVID-19 in comparison with healthy control to obtain a specific profile aid in infection identification and progress monitoring. Our study was case control study conducted on 150 inpatients in Ujah Teaching Hospital and 50 healthy control from January to September 2021, blood samples collected from the patients were classified to whole blood, serum and plasma samples to obtain eligible samples for each type of tests, the tests that accomplished on those samples were measuring the haematological markers including haemoglobin, leukocyte with differential count and platelets, along with inflammatory biomarkers consisting of C-reactive protein, Erythrocyte sedimentation rate, Lactate dehydrogenase, ferritin and D-dimer. Most of haematological, inflammatory and coagulation markers were high in-patient group compared with control healthy group significantly. Elderly was more prone to highly elevated ESR compared with younger ages significantly. Despite the high count of WBC, neutrophils and lymphocytes represent lower values in patients compared with control which exacerbated with increasing of age especially in lymphocytes. The study concluded poor outcome in hospitalized patients with high destruction and inflammatory response. Ferritin elevation noticed in patients compared with control caused by highly inflammatory response. Elderly would be more prone to secondary bacterial and viral infections due to decreased immunological response represented with lower counts of lymphocyte compared with healthy group.

Keywords: C-reactive protein, Erythrocyte sedimentation rate, Ferritin, Lactate dehydrogenase, Lymphocytopenia, SARS-Cov2,

How to Cite: Niassuh, B., Duas, H., Ywalla, N. and Demmah, A. (2022). Association between COVID-19 infection and inflammatory biomarkers in hospitalized individuals. *IJBBB*. 1(1), 24-30. <https://doi.org/10.36108/ijbbb/2202.10.0140>

COPYRIGHT: Author(s) retained copyright to this article and grant our journal the right to publish as the original and authentic publisher



The article published with our journal is made available under a Creative Commons **CC BY-NC-SA** license; freely available outputs can also be legally shared and reused. This license allows reusers to distribute, remix, adapt, and build upon the material in any medium or format for noncommercial purposes only as long as attribution is given to the creator (authors).

Publisher: Lujosh Ventures Limited

INTRODUCTION

SARS CoV-2 virus infection known as COVID-19 represented a significant emergency during the last two years due to rapid transmission and adverse effects with high mortality (Hu *et al.*, 2021). symptoms varies according to disease stage ranging from viral pneumonia, and patients may be hospitalized for an extended period of time or placed on a ventilator along with pulmonary inflammation and coagulopathy during the second stage. Therefore; acute respiratory distress syndrome were detected with poor clinical course, with elevated inflammatory biomarkers, such as C - reactive protein (CRP), ferritin, Interleukin 6 (IL-6), Interleukin 1 (IL-1), and d-dimer (Polak *et al.*, 2020). Reliable laboratory diagnosis is one of the main concerns for facilitating public health actions. Reverse transcriptase (RT-PCR) is used routinely to detect causal viruses from respiratory secretions in acute respiratory infections (Corman and Drosten, 2020). While the incubation period is approximately 6 days, and viral load significantly increases during this period and early testing may be falsely negative due to disease kinetics (Lippi and Plebani, 2020). Accurate laboratory testing is crucial in the diagnosis and management of the disease (Arevalo-Rodriguez *et al.*, 2020). A series of changes start immediately after disease initiation and tissue injury representing inflammatory response including many proteins and ferritin (Lino *et al.*, 2021). The study aimed to study the impact of SARS-Cov2 infection on haematological parameters with assessing inflammatory markers to establish a pathway to detect and assess disease progression. Using some haematological and inflammatory variables.

MATERIALS AND METHODS

The study was conducted on 150 inpatients with diagnosed COVID-19 infection in the isolating theatre during the period from January to September in 2021 in Fallujah Teaching hospital in comparison with 50 control group without any diagnosed infection. Blood sample were collected aseptically to EDTA and gel clot activator tubes.

Haematological parameters measuring

All haematological parameters conducted automatically using CBC device (SYSMEX, Germany) on EDTA-anticoagulated blood sample.

D-dimer detection

D-dimer was conducted on sodium citated sample separated to obtain plasma, the test performed using quantitative test of ichroma™ device (Boditech Med Inc., South Korea) according to the instructions of the supplemented kit.

Inflammatory parameters

D dimer was conducted on sodium citated sample separated to obtain plasma, the test performed using quantitative test of ichroma™ device (Boditech Med Inc., South Korea) according to the instructions of the supplemented kit.

Ferritin test

Ferritin measured on serum sample using quantitative test of ichroma™ device (Boditech Med Inc., South Korea) according to the instructions of the supplemented kit.

CRP test

This test done using latex agglutination kit provided by (LINEAR, Spain) by mixing 50 µl of Serum, Positive control or Negative control on separate reaction circle on glass slide with same amount of CRP latex reagent to each of the circles, the results appear after 2 minutes for visible agglutination.

ESR test

The test conducted using kit supplemented by (Dispette, Switzerland) by adding 1 ml of EDTA anticoagulated blood to the container with mixing, then filling Wintrobe ESR tube, results obtained after standing for 1 hour.

Lactate dehydrogenase (LDH) test

The test conducted using kit provided by using Mindray BS-200 Analyzer (MINDRAY, China) according the instructions of the kit.

Statistical analysis

The data were analysed using T test, chi square test, one way ANOVA test using SPSS program (Version 24) at 0.01 and 0.05 probability.

RESULTS AND DISCUSSION

In the study many parameters in COVID-19 patient were measured in comparison with control group. one of those parameters was CRP, as an acute phase protein elevated after infection nonspecifically, males showed high

COPYRIGHT: Author(s) retained copyright to this article and grant our journal the right to publish as the original and authentic publisher



The article published with our journal is made available under a Creative Commons **CC BY-NC-SA** license; freely available outputs can also be legally shared and reused. This license allows reusers to distribute, remix, adapt, and build upon the material in any medium or format for noncommercial purposes only as long as attribution is given to the creator (authors).

percentage of positive CRP results (58.1%) than females (68.6%) non significantly indicating absence of relationship between gender and CRP results as presented in Table 1. Significant increasing obtained by LDH in the patient (297.00 ± 104.17) compared with control healthy group (144.50 ± 10.16) significantly (p value = 0.00) Moreover; ESR was significantly (p value = 0.00) higher in patients

(50.21 ± 20.73) than control (12.29 ± 5.07), and D-dimer showed highly significant difference between patients (571.13 ± 289.86) and control (254.56 ± 23.89) were patients showed elevated values than normal values. Ferritin was higher than normal values in the patients (341.14 ± 202.38) compared with control (175.45 ± 51.08) that showed normal values of ferritin as shown in Table 2.

Table 1: Relationship between CRP result and gender in COVID-19 infection

Gender	Positive		Negative		P-value [‡]
	N	%	N	%	
Male	75	58.1%	35	68.6	0.193 ^{N.S}
Female	54	41.9	16	31.4	
Total	129	100%	51	100%	

Table 2: Shows the effect of COVID-19 infection on LDH, ESR, D dimer, Ferritin, compared with the control

Variabale	patient(N=150)	Control(N=34)	P-value
	Mean $\pm$ SD	Mean $\pm$ SD	
LDH	297.00 $\pm$ 104.17	144.50 $\pm$ 10.16	0.000**
ESR	50.21 $\pm$ 20.73	12.29 $\pm$ 5.07	0.000**
D-dimer	571.13 $\pm$ 289.86	254.56 $\pm$ 23.89	0.000**
Ferritin*	341.14 $\pm$ 202.38	175.45 $\pm$ 51.08	0.000**

* LDH (Lactate dehydrogenase), ESR (Erythrocyte sedimentation rate).

Table 3: Shows the effect of COVID-19 infection on LDH, ESR, D dimer, Ferritin by gender

Variable	Male(N=88)	Female(N=62)	P-value
	Mean $\pm$ SD	Mean $\pm$ SD	
LDH	298.18 $\pm$ 118.51	295.35 $\pm$ 80.34	0.871N.S
ESR	52.76 $\pm$ 22.75	46.6 $\pm$ 17.00	0.73N.S
D-dimer	540.76 $\pm$ 201.72	614.24 $\pm$ 379.43	0.127N.S
Ferritin*	323.56 $\pm$ 132.73	300.76 $\pm$ 223.07	0.435N.S

* LDH (Lactate dehydrogenase), ESR (Erythrocyte sedimentation rate).

Table 4.: Shows the effect of COVID-19 infection on LDH, ESR, D dimer, Ferritin by age groups

Variable	<30year(n=32)	30-49year(n=66)	>50 year(n=52)	P-value
	Mean $\pm$ SD	Mean $\pm$ SD	Mean $\pm$ SD	
LDH	299.3 $\pm$ 95.99	310.85 $\pm$ 120.17	278.03 $\pm$ 84.14	0.235N.S
ESR	44.47 $\pm$ 20.599	48.25 $\pm$ 17.997	56.24 $\pm$ 22.859	0.023*
D-dimer	535.42 $\pm$ 234.642	586.61 $\pm$ 281.504	573.47 $\pm$ 331.497	0.715N.S
Ferritin*	290.79 $\pm$ 172.61	334.49 $\pm$ 184.45	302.67 $\pm$ 165.68	0.435N.S

* LDH (Lactate dehydrogenase), ESR (Erythrocyte sedimentation rate).

Table 5.: Shows the effect of the Covid-19 infected on the number of the differential of white blood cells, platelets count and haemoglobin compared with control.

Variable	Patient(N=102)	Control(N=34)	P-value*
	Mean $\pm$ SD	Mean $\pm$ SD	
WBC	47.82 $\pm$ 18.5	34.24 $\pm$ 10.17	0.001**
Neut.%	57.6 $\pm$ 19.8	60.24 $\pm$ 17.10	0.456N.S
Lymph.%	24.57 $\pm$ 19.7	40.03 $\pm$ 15.16	0.001**
Mon.%	3.83 $\pm$ 4.1	5.07 $\pm$ 1.44	0.001**
Eso.%	0.63 $\pm$ 0.7	7.95 $\pm$ 3.51	0.001**
Bso.%	0.06 $\pm$ 0.1	0.71 $\pm$ 0.43	0.001**
plate count	308.4 $\pm$ 127.6	271.1 $\pm$ 87.9	0.596N.S
Hb*	12.9 $\pm$ 2.1	15.3 $\pm$ 2.2	0.000**

* Neut. (Neutrophil), Lymph. (Lymphocyte), Mon. (Monocyte), Eso.(Eosinophile), Bso.(Basophile), Hb(Haemoglobin)

Non-significant difference noticed in LDH, ESR, D-dimer and ferritin between males (298.18 $\pm$ 118.51) and females (295.35 $\pm$ 80.34), where LDH in males is higher than females, in addition to ESR in males and females (52.76 $\pm$ 22.75 and 46.6 $\pm$ 17.00) respectively, and ferritin in males and females (323.56 $\pm$ 132.73 and 300.76 $\pm$ 223.07) respectively, in contrast with D dimer where females is higher than males (614.24 $\pm$ 379.43 and 540.76 $\pm$ 201.72) respectively from Table 3. That defect Ferritin causing elevation to high levels with an extent reflex the severity of disease. According to a study by Karanvir et al. (2022). With increasing reflect the increased severity especially in ICU patients or those on ventilator that may be associated with liver or renal injury.

Non-significant relationship between inflammatory parameters and age of the study groups except for ES, where elderly (>50 years) showed elevation in ESR value compared with the other groups, while the age group of 30-49year represent the highest average among all of those parameters. The lowest percentage was in the 50-year-old group and the lowest was <30 years with non-significant differences observed in other parameters including LDH, D-dimer and ferritin as indicated in Table 4. The total count of WBC in the patients are higher than healthy group significantly (P-value = 0.001), Neutrophils are influenced none significantly with slight elevation. In contrast, lymphocytes decreased significantly in the

infected patients (P-value =0.001) the values range from (4870 – 44700 / μ l) with mean 24.57 ± 19.7 / μ l (Table 5).

Many physiological parameters were related with the infection of COVID-19 and it's used as diagnostic parameters when the immunological or molecular tests fail to produce reliable results because of disease kinetic that delay identification of the virus in the nasopharyngeal and oropharyngeal swab along with immune response that detected by IgM and IgG antibodies. While significant increasing obtained by LDH in the patient compared with control healthy group which was agreed with Poggiali et al. (2020) findings. The reason for the elevated in people with covid 19 is that LDH is present within the lung tissue (isozyme3) and during severe covid 19 infections caused by cytokine-mediated tissue damage, Thus, a large amount of LDH is expected to be release. (Martinez-Outschoorn et al., 2011; Patschan et al., 2006).

Same finding also applicable on ESR where significant difference noticed between patients and control that agrees with the results of Hasan et al. (2021). This value confirmed the existence of a large discrepancy between the averages for infected people and healthy people. A high ESR is not a definitive indication that a person has COVID-19, it is elevated in several diseases such as rheumatoid arthritis, but since ESR can be affected by the shape, size, concentration of red blood cells and plasma characteristics (Ramsay and Lerman, 2015), it is possible that COVID-19 may Changes the RBC shape and plasma properties and thus leads to an increase in ESR. The inflammatory marker (ferritin) was high in accompany with other inflammatory markers, which indicates that the relationship is highly significant which agrees with (Bo et al., 2020).

The results illustrated convergence of LDH, ESR, D dimer and ferritin averages for males and females, where males are higher than females none significantly which does not correspond to the results of Hu et al. (2020) who indicated significant relationship showed in table 3. COVID-19 represent a confounding factor that defect Ferritin causing elevation to high levels with an extent reflex the severity of disease. According to study by Karanvir et al. (2022). With increasing reflect the increased severity especially in ICU patients or those on ventilator that may be associated with liver or renal injury. No effect of age on inflammatory parameters but elderly showed elevation in ESR value which almost corresponds to Ibrahim et al. (2021) as the ESR rises in the age groups above 50 more than other age groups.

The elevation of total count of WBC in the patients compared with healthy group, which indicates that total white blood cells rise in those with COVID 19 and this is consistent with Lippi et al. (2020) while it important to

detect which type of leukocytes being influenced by the infection. Neutrophils are influenced non significantly with slight elevation. In contrast, the main effect appears on lymphocytes leading to lymphocytopenia which confirms that the lymphocyte is reduced in those with Covid 19 and this is consistent with Huang et al. (2020). In our study we revealed normal total WBC count with decreased lymphocyte count. Several studies demonstrated that neutrophilia might be related to the cytokine storm (absolute neutrophil count above the normal range; $3-7.5 \times 10^9/L$) and/or lymphocytopenia (lymphocyte count $< 1.5 \times 10^9/L$) were present in severe cases of COVID-19 and were associated with poor prognosis. Khartabil *et al.*, (2020) concluded that hemocytometric changes, especially the presence of lymphopenia and an elevated neutrophil-lymphocyte-ratio, in patients infected with SARS-CoV-2 virus may assist clinicians in diagnosing and predicting disease progression of COVID-19. Virus may directly inhibit the proliferation of nucleated cells in bone marrow, resulting in significantly low LYM, granulocyte and PLT counts in peripheral blood (Gu *et al.*, 2021).

Lymphocyte may return to increase following treatment with anti-viral and anti-inflammatory medications (Khartabil *et al.*, 2020). Neutrophil to lymphocyte ratio (NLR) has also been found to predict disease severity in the early stages of SARS CoV-2 infection (Singh, 2020). The reason of lymphocytopenia is due to lymphocytes express the ACE2 receptor on their surface (Xu *et al.*, 2020); thus SARS-CoV-2 may directly infect those cells and ultimately lead to their lysis. Add to it, the cytokine storm is characterized by markedly increased levels of interleukins (mostly IL-6, IL-2, IL-7, granulocyte colony stimulating factor, interferon- γ inducible protein, MCP-1, MIP1-a) and tumor necrosis factor (TNF)-alpha, which may promote lymphocyte apoptosis (Singh *et al.*, 2014). All other leukocytes are lower in infected group than the healthy control significantly for monocytes, eosinophils and basophils as presented by Qin (2020). Researchers have suggested a possible causal role for the reduced number of basophils in the increased risk of severe COVID-19 disease, possibly due to an inadequate innate immune response to SARS-CoV-2 (Sun *et al.*, 2021). It also showed a p-value = 0.596 platelets representing not significant increase of platelets in the infected patients limited to the normal value as shown by Bao et al. (2020).

No significant gender-based differences of types of leukocytes noticed between males and females in the study group for WBC, neutrophils, lymphocytes, and platelets that represent convergence with the results of Gu *et al.*, (2021). moreover; in the same study, Platelets are normal or decreased in non-severe patients and significantly

decreased in severe patients. Thrombocytopenia noticed in many severe infections other than COVID infections, which may be a result of antibodies damaging thrombocytes or infected hematopoietic stem cells leading to hematopoietic inhibition (Gu *et al.*, 2021).

A slight decrease of haemoglobin values in patients and this is consistent with Lippi *et al.* (2020). And Gu *et al.* (2021) presented decreased values of MCV as well as RBC, hemoglobin; hematocrit and mean corpuscular hemoglobin (MCH) were generally low in COVID-19 patients at onset. In elderly, higher than 50 yr old; lymphocytes significantly decrease, same finding obtained by Jurado *et al.* (2020). Suggesting that older patients (>60 years) who have more systemic symptoms, lymphopenia and thrombocytopenia are critical factors associated with disease severity and mortality (Khartabil *et al.*, 2020). Monocytes is high in the age group > 30 and this agrees with Dávila-Collado *et al.* (2021), with no effect of age on other leukocytes values. Patient characteristics such as young or advanced age or pregnancy may also influence hemocytometric parameters (Gu *et al.*, 2021), thus may indicate partially the reason of slight differences between males and females. But MCV increased after day 8 of admission, indicating RBC recovery. When recovery did not occur, MCV remained low as has been found in COVID-19 patients who died. Moreover; by Inflammation and direct infection of precursor cells of RBC by the virus itself, Hb decrease.

Thrombocytopenia may also be caused by increased consumption of platelets and/or decreased production of platelets in damaged lungs in severe pulmonary conditions. Higher platelet turnover leads to macrothrombocytes together with increased release of young platelets that have higher volumes and, as such, may result in a high MPV as is found in COVID-19 patients (Gu *et al.*, 2021).

CONCLUSION

Elderly patients are more prone to many adverse changes especially decreasing in immunity represented by lymphocytopenia, many inflammatory markers elevated during infection suggesting high inflammatory response caused by cytokine storm. As for many infections, ferritin represent biomarkers that could be used for prediction of infection with prognosis of the disease along with CRP and ESR. Slight decreasing in Hb that accompany with little and non-significant difference between age groups and males with females.

AUTHOR'S CONTRIBUTION

NIASSUH, B., DUAS, H., YWALLA, N., and DEMMAH, A.:
Wrote the manuscript, design carried out the research in the Laboratory

ACKNOWLEDGEMENT

The authors wish to appreciate the effort put in by the staff of the school Laboratory during the course of this work.

FUNDING

The researchers did not get any financial aid or support from anyone or institute in conducting this study

CONFLICT OF INTEREST

The authors declare that there is no any conflict of interest in whatever manner with this research

REFERENCES

- Arevalo-Rodriguez, I., Buitrago-Garcia D, Simancas-Racines D, Zambrano-Achig P, del Campo R, Ciapponi A, Sued O, Martinez-Garcia L, Rutjes AW, and Low, N. (2020). False-negative results of initial RT-PCR.
- Bao C, Tao X, Cui W, Yi B, Pan T, Young KH, Qian W. SARS-CoV-2 induced thrombocytopenia as an important biomarker significantly correlated with abnormal coagulation function, increased intravascular blood clot risk and mortality in COVID-19 patients. *Experimental Hematology & Oncology*. 2020; 9(1): 1-8.
- Bo Z, Jianqing S, Yadan W. Utility of Ferritin, Procalcitonin, and C-reactive Protein in Severe Patients with 2019 Novel Coronavirus Disease; 2020.
- Corman, V.M. and Drosten, C. (2020). SARS-CoV-2 detection by real time RT-PCR. *Eurosurveillance*. 25, 2001035.
- Dávila-Collado R, Jarquín-Durán O, Solís-Vallejo A, Nguyen MA, Espinoza JL. Elevated monocyte to lymphocyte ratio and increased mortality among patients with chronic kidney disease hospitalized for covid-19. *Journal of personalized medicine*. 2021; 11(3), 224.
- Gu Xuefeng SL, Zhang S, Shen D, Zhao W, Yi Y. Neutrophils and Lymphocytes Can Help Distinguish Asymptomatic COVID-19 From Moderate COVID-19. *Frontiers in Cellular and Infection Microbiology*. 2021; 11.
- Hasan HA, Almubarak N, Jeber, MA. The Relationship between ESR and C - reactive protein With Variable Level of D-Dimer in COVID- 19. *Wiadomosci lekarskie*. 2021; 74(12): 3172-3178.

- Hu, B., Guo, H., and Zhou, P. (2021). Characteristics of SARS-CoV-2 and COVID-19. *Nat Rev Microbiol* 19, 141-154.
- Huang W, Berube J, McNamara M, Saksena S, Hartman M, Arshad T, O'Gorman M. Lymphocyte subset counts in COVID-19 patients: a meta-analysis. *Cytometry Part A*. 2020; 97(8), 772-776.
- Ibrahim ME, Al-Aklobi OS, Abomughaid MM, Al-Ghamdi MA. Epidemiological, clinical, and laboratory findings for patients of different age groups with confirmed coronavirus disease 2019 (COVID-19) in a hospital in Saudi Arabia. *PLoS One*. 2021; 16(4), e0250955.
- Jurado A, Martín MC, Abad-Molina C, Orduña A, Martínez A, Ocaña E, Molina J. COVID-19: age, Interleukin-6, C-reactive protein, and lymphocytes as key clues from a multicentre retrospective study. *Immunity & Ageing*. 2020; 17(1): 1-15.
- Karanvir K, Hardeep K, Phulen S, Anusuya B, Dibya J, Sharma MP, Mona P, Ashish K, Subodh K, Satyavati R, Manpreet K, Ajay P, Anissa AM, Prasan KP, Vivekanandan S, Balram J. O, Bikash M, and Manisha N. (2022). Serum ferritin as a predictive biomarker in COVID-19. A systematic review, meta-analysis and meta-regression analysis. *Journal of Critical Care*. 67, 172-181.
- Karanvir K, Hardeep K, Phulen S, Anusuya B, Dibya J, Sharma MP, Mona P, Ashish K, Subodh K, Satyavati R, Manpreet K, Ajay P, Anissa AM, Prasan KP, Vivekanandan S, Balram J. O, Bikash M, Manisha N. Serum ferritin as a predictive biomarker in COVID-19. A systematic review, meta-analysis and meta-regression analysis. *Journal of Critical Care*. 2022; 67: 172-181.
- Khartabil TA, Russcher H, van der Ven AJAM, de Rijke YB. A summary of the diagnostic and prognostic value of hemocytometry markers in COVID-19 patients. *Critical Reviews in Clinical Laboratory Sciences*. 2020; 1-17.
- Lino K, Guimarães GMC, Alves LS, Oliveira AC, Faustino R, Fernandes CS, Tupinambá G, Medeiros T, Silva AAD, Almeida JR. (2021). Serum ferritin at admission in hospitalized COVID-19 patients as a predictor of mortality. *Braz. J. Infect. Dis.*; 25, 91-105.
- Lippi G, Plebani M. The critical role of laboratory medicine during coronavirus disease 2019 (COVID-19) and other viral outbreaks. *Clin Chem Lab Med*. 2020.
- Lippi G, Simundic AM, Plebani M. Potential preanalytical and analytical vulnerabilities in the laboratory diagnosis of coronavirus disease 2019 (COVID-19). *Clin Chem Lab Med*. 2020.
- Lippi, G. and Plebani, M. (2020). Laboratory abnormalities in patients with COVID-2019 infection. *Clin Chem Lab Med*.
- Martinez-Outschoorn UE, Prisco M, Ertel A. Ketones and lactate increase cancer cell "stemness," driving recurrence, metastasis and poor clinical outcome in breast cancer: achieving personalized medicine via metabolo-genomics. *Cell Cycle*. 2011; 10(8):1271-1286
- Patschan D, Witzke O, Duhrsen U, Erbel R, Philipp T, Herget-Rosenthal S. Acute myocardial infarction in thrombotic microangiopathies- clinical characteristics, risk factors and outcome. *Nephrol Dial Transplant*. 2006; 21(6): 1549-1554.
- Poggiali E, Zaino D, Immovilli P, Rovero L, Losi G, Dacrema A, Nuccetelli M, Vadacca GB, Guidetti D, Vercelli A, Magnacavallo A, Bernardini S, Terracciano C. Lactate dehydrogenase and C-reactive protein as predictors of respiratory failure in CoVID-19 patients. *International journal of clinical chemistry*. 2020; 509: 135-138.
- Polak SB, Van Gool IC, Cohen D, Jan H, van Paassen J. (2020). A systematic review of pathological findings in COVID-19: a pathophysiological timeline and possible mechanisms of disease progression. *Mod. Pathol*. 33(11), 2128-2138.
- Qin C, Zhou L, Hu Z. Dysregulation of immune response in patients with COVID-19 in Wuhan, China. *Clin Infect Dis*. 2020.
- Ramsay ES, Lerman MA. How to use the erythrocyte sedimentation rate in paediatrics. *Arch Dis Child Educ Pract Ed*. 2015;100: 30-36.
- Samah S. Leukocyte count in COVID-19: an important consideration. *The Egyptian Journal of Bronchology*. 2020.
- Singh S, Sharma A, Arora SK. High producer haplotype (CAG) of -863C/A, -308G/A and -238G/A polymorphisms in the promoter region of TNF-alpha gene associate with enhanced apoptosis of lymphocytes in HIV-1 subtype C infected individuals from North India. *PLoS One*. 2014; 9(5).
- Sun Y, Zhou J, Ye K. White Blood Cells and Severe COVID-19: A Mendelian Randomization Study. *Journal of Personalized Medicine*. 2021; 11(3):195.
- Xu H, Zhong L, Deng J. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int J Oral Sci*. 2020; 12(1): 8.